

## Baljit Dheansa FRCS MSc FRCS(Plast) Consultant Plastic, Cosmetic & Reconstructive Surgeon

Му	Plastic Surgeon Limited
	The McIndoe Centre
	Holtye Road
	East Grinstead
	West Sussex
	RH19 3FR

Patient: Procedure	э:	Gynaecomastia On Date:			
	•				
RISKS / COMPLICATIONS (rare or unusual complications can occur and may not be listed) General					
Allergic reactions to any drugs, antiseptics	or	Dehiscence – wound opening up – may require			
equipment used	-	prolonged dressings			
Asymmetry (this may be better or worse than p	ore-				
operatively)					
Bleeding – may require further procedure		dressings or surgery including skin grafts  Swelling, Inflammation – may take months to			
bleeding – may require further procedure		resolve			
Bruising – may take weeks to resolve	Bruising may take weeks to resolve				
Haematoma (collection of blood under skin) – r	nav	Slow healing Poor scarring – Keloid, Hypertrophic, unsightly –			
require further procedure	iiay	may be difficult to treat or improve			
Infection – may require antibiotics or surgery or		may increase risk of inadvertent injury in future			
may cause tissue loss		Pulmonary Embolus or Deep Vein Thrombosis –			
Pain – may be long term					
Changes in contour of tipours, may be view	، داله د	clots in the lung or legs			
Changes in contour of tissues – may be visu	ally				
apparent		droop or lose the effect of surgery			
Specific to: Gynaecomastia					
Change in shape of areola - may not be rour	nd,	Scars that go round the whole areola may be ruched			
may remain large, maybe become indented	or	and may remain so			
parts of it may be less obvious					
Dishing of the areola – inward position which may		Scars may be visible especially if need to be			
be difficult to treat	-	extended.			
Seroma – fluid collection		Residual lumpiness			
Tethering or dimpling of the skin					
Incomplete removal, not enough removed		Further scars around the chest if having liposuction  May require a second stage or further surgery			
Over correction, too much removed		Disappointment with final result			
Loss of effect over time or with weight gain		Stretch marks may remain			
Loss of the nipple - partial or total - may	be	Skin may become loose afterward and not retract			
difficult to reconstruct		fully			
Mr Baljit Dheansa has discussed the above risks and complications with me. I understand these. I have read Mr Dheansa's Gynaecomastia information leaflet [ (Please Tick)]. I also understand other risks and complications can occur but may be less common as the practice of medicine and surgery is not an exact science. I understand the risks, benefits and limitations of the proposed procedure. Certain complications may affect my ability to work, socialise or do certain activities. This may sometimes be long term.  I permit the use of my non-identifiable photographs (pre-and post-operative) to be used for teaching, publication and to be shown to other patients. I can withdraw this permission at any time.  Patient Signature:					
Patient Name:		Date:			

Mr BS DHEANSA FRCS MSc FRCS(Plast) Consultant Plastic, Reconstructive & Cosmetic Surgeon

Telephone: 01342 330 383 Facsimile: 01342 330 301

Email: <a href="mailto:enquiry@my-plastic-surgeon.co.uk">enquiry@my-plastic-surgeon.co.uk</a> Website: <a href="mailto:www.my-plastic-surgeon.co.uk">www.my-plastic-surgeon.co.uk</a>



## **ADDITIONAL NOTES OR COMMENTS:**

ı	
ı	
ı	
ı	
ı	
ı	
ı	
ı	
ı	
ı	
ı	
ı	
ı	
ı	
ı	
ı	
ı	
ı	
ı	
ı	
ı	
ı	
1	

Telephone: 01342 330 383 Facsimile: 01342 330 301 Email: <a href="mailto:enquiry@my-plastic-surgeon.co.uk">enquiry@my-plastic-surgeon.co.uk</a> Website: <a href="mailto:www.my-plastic-surgeon.co.uk">www.my-plastic-surgeon.co.uk</a> Website: www.my-plastic-surgeon.co.uk